



THE CAUSES OF TYPE 2 DIABETES

**STATISTICS, RISK FACTORS, PATHOPHYSIOLOGY AND AN
INTEGRATED HOLISTIC WELLNESS APPROACH**

AN INTEGRATIVE CLINICAL RESEARCH REVIEW

Prepared by

A. SUDHAKAR

Qualified Ayurvedic Therapist & Panchakarma Specialist

**SR VAIDYA HOLISTIC WELLNESS RESEARCH INSTITUTE
SRVAIDYA.com**

Understanding Type 2 Diabetes Beyond Blood Sugar: A Multifactorial Metabolic Disease

ABSTRACT

Type 2 Diabetes Mellitus (T2DM) is a chronic metabolic disease characterized by persistent hyperglycemia arising primarily from insulin resistance and progressive impairment of pancreatic beta-cell function. It is one of the major global public-health challenges and should not be reduced to the oversimplified idea that it is caused by sugar alone. Its development reflects the interaction of genetic susceptibility, excess adiposity, physical inactivity, dietary patterns, aging, metabolic dysfunction and other environmental influences.

This review examines the principal causes and risk factors associated with T2DM, including insulin resistance, visceral adiposity, beta-cell dysfunction, prediabetes, metabolic syndrome, fatty liver disease, family history, gestational diabetes, polycystic ovary syndrome, sleep and lifestyle influences. It also discusses evidence-based prevention and proposes an HWRI–T2D Integrated Care Model emphasizing screening, assessment, identification of risk factors, correction, optimization, monitoring and empowerment.

Keywords: Type 2 Diabetes, insulin resistance, beta-cell dysfunction, obesity, visceral fat, prediabetes, metabolic syndrome, lifestyle medicine, diabetes prevention, metabolic health, holistic wellness.

1. INTRODUCTION

Diabetes occurs when the body does not produce enough insulin, cannot effectively use the insulin it produces, or both. In Type 2 Diabetes, the dominant processes are reduced insulin sensitivity and progressive failure of insulin secretion to compensate adequately. The disease may evolve gradually over many years and can remain clinically silent during its early metabolic stages.

TYPE 2 DIABETES IS NOT SIMPLY A “SUGAR DISEASE.” IT IS A PROGRESSIVE DISORDER OF METABOLIC REGULATION.

2. GLOBAL BURDEN AND STATISTICS

The World Health Organization reports that approximately 830 million people were living with diabetes globally in 2022, representing a substantial rise compared with 1990. Type 2 Diabetes accounts for the majority of diabetes cases. The burden extends beyond elevated glucose because persistent dysglycemia is associated with cardiovascular disease, stroke, chronic kidney disease, retinopathy, neuropathy and lower-limb complications.

Indicator	Significance
Global burden	Approximately 830 million people living with diabetes in 2022
Trend	Major increase since 1990
Major mechanisms	Insulin resistance + progressive beta-cell dysfunction
Major complications	Heart, brain, kidney, eye, nerve and vascular damage

3. NORMAL GLUCOSE–INSULIN PHYSIOLOGY

After food is digested, glucose enters the bloodstream. The pancreas releases insulin, which helps insulin-responsive tissues such as skeletal muscle and adipose tissue take up glucose, while also regulating hepatic glucose production. Glucose is then used for energy or stored, helping maintain blood glucose within a physiological range.

FIGURE 1. Normal Glucose–Insulin Physiology



Normal glucose regulation: food-derived glucose enters the bloodstream, insulin is released by the pancreas, and glucose is taken up by responsive tissues for energy or storage.

4. WHAT GOES WRONG IN TYPE 2 DIABETES?


A common disease sequence involves insulin resistance followed by compensatory increases in insulin secretion. Over time, pancreatic beta cells experience metabolic stress and may no longer produce enough insulin to maintain normal glucose levels. The result is progressive dysglycemia and eventually persistent hyperglycemia.

INSULIN RESISTANCE → COMPENSATORY HYPERINSULINEMIA → BETA-CELL STRESS → RELATIVE INSULIN DEFICIENCY → PERSISTENT HYPERGLYCEMIA → TYPE 2 DIABETES

5. INSULIN RESISTANCE — THE CENTRAL MECHANISM

Insulin resistance means that target tissues respond less effectively to insulin. In skeletal muscle, glucose uptake is reduced; in the liver, insulin may inadequately suppress glucose production; and dysfunctional adipose tissue may contribute to altered fatty-acid metabolism and inflammatory signaling. These processes interact and can accelerate metabolic deterioration.

FIGURE 3. Insulin Resistance: Normal vs Impaired Glucose Uptake




Comparison of normal insulin action with impaired cellular glucose uptake, showing the progression toward hyperglycemia and beta-cell stress.

6. OVERWEIGHT, OBESITY AND VISCERAL FAT

Excess body fat, particularly abdominal or visceral adiposity, is an important risk factor for Type 2 Diabetes. Visceral adipose tissue is metabolically active and may contribute to chronic low-grade inflammation, altered lipid flux and insulin resistance. However, Type 2 Diabetes can also occur in people without obesity; genetic susceptibility, ectopic fat distribution and other metabolic factors may be important.

FIGURE 4. Visceral Fat → Insulin Resistance → Type 2 Diabetes



A metabolic pathway illustrating how excess visceral adiposity contributes to inflammatory and lipid-related dysfunction and increased diabetes risk.

7. PHYSICAL INACTIVITY

Skeletal muscle is a major site of glucose disposal. Physical inactivity may reduce insulin sensitivity, decrease energy expenditure and contribute to weight gain and worsening metabolic health. Regular aerobic and resistance activity can improve insulin sensitivity and is a key component of diabetes prevention strategies.


8. DIETARY PATTERNS AND TYPE 2 DIABETES

It is scientifically incomplete to state that eating sugar alone causes Type 2 Diabetes. Dietary patterns can influence total energy intake, body weight, visceral adiposity, insulin sensitivity and overall cardiometabolic health. Frequent intake of highly processed foods, low fibre intake and chronic excess calorie consumption may contribute to metabolic risk.

9. GENETICS AND FAMILY HISTORY

Family history and genetic susceptibility are important non-modifiable contributors. Yet genetic risk is not destiny: genes interact with lifestyle, body composition, physical activity, dietary patterns, age and environmental exposures.

FIGURE 2. Comprehensive Causes of Type 2 Diabetes



A multifactorial overview of the interacting metabolic, genetic, lifestyle and environmental contributors associated with Type 2 Diabetes.


10. AGE AND TYPE 2 DIABETES

Risk generally rises with age because of changes in body composition, reduced physical activity, progressive insulin resistance and declining beta-cell reserve. Nevertheless, Type 2 Diabetes increasingly affects younger adults and can also occur in adolescents and children.

11. PREDIABETES — THE WARNING STAGE

Prediabetes describes glucose levels above normal but below diagnostic thresholds for diabetes. It often represents an important opportunity for early detection and risk reduction because progression is not inevitable and lifestyle intervention may delay or prevent Type 2 Diabetes in high-risk individuals.

FIGURE 5. The Progression Pathway: Normal Metabolism → Prediabetes → Type 2 Diabetes



Poor sleep patterns and sleep disorders can influence appetite regulation, body weight, hormonal balance and insulin sensitivity. Sleep should be viewed as one component of a broader metabolic and lifestyle network rather than as an isolated cause.


17. STRESS AND METABOLIC HEALTH

Chronic stress may influence eating behaviour, physical activity, sleep and neuroendocrine pathways. Stress management can support overall wellbeing, but stress alone should not be presented as a single cause of Type 2 Diabetes.

18. SMOKING, MEDICATIONS AND OTHER CONTRIBUTORS

Smoking is associated with increased metabolic and cardiovascular risk. Certain medications may also alter glucose metabolism in susceptible individuals. Medication-related hyperglycemia requires appropriate clinical evaluation rather than automatic classification as typical Type 2 Diabetes.

FIGURE 6. Lifestyle Risk Factor Pyramid



A visual summary of modifiable lifestyle influences including dietary patterns, physical inactivity, sleep, stress and other metabolic risks.

19. THE SILENT PROGRESSION OF TYPE 2 DIABETES

Many individuals pass through stages of insulin resistance, compensatory hyperinsulinemia and prediabetes before clinical diagnosis. Because this process may occur without obvious symptoms, risk-based screening and early detection are essential.

20. COMPLICATIONS OF PERSISTENT HYPERGLYCEMIA

Persistent hyperglycemia can damage blood vessels and nerves, contributing to cardiovascular disease, stroke, diabetic kidney disease, retinopathy and vision loss, peripheral neuropathy and foot complications. Prevention of complications requires comprehensive risk-factor management, not glucose control alone.


21. EVIDENCE-BASED PREVENTION

Key prevention strategies include healthy dietary patterns, regular physical activity, sustainable weight management where appropriate, smoking cessation, early screening and management of prediabetes. Lifestyle interventions should be individualized and integrated with qualified medical care.

22. THE ROLE OF HOLISTIC WELLNESS

A holistic approach can support comprehensive care by integrating medical assessment, nutrition, physical activity, weight management, sleep optimization, stress management and patient education. Such approaches should complement—not replace—diagnosis, prescribed medication or evidence-based diabetes management.

FIGURE 7. Ayurvedic Perspective: Prameha (Madhumeha) & Modern Correlation



An integrative conceptual discussion of Prameha/Madhumeha, Ayurvedic principles and their supportive correlation with modern metabolic understanding.

23. HWRI–T2D INTEGRATED CARE MODEL™

SCREEN → ASSESS → IDENTIFY → CORRECT → OPTIMIZE → MONITOR → EMPOWER

SCREEN: identify people at increased risk. **ASSESS:** evaluate glucose, HbA1c, body composition, blood pressure, lipids and lifestyle patterns. **IDENTIFY:** recognize major contributing metabolic and behavioural factors. **CORRECT:** address modifiable risks and ensure appropriate medical care. **OPTIMIZE:** improve nutrition, activity, sleep and metabolic health. **MONITOR:** follow clinically relevant indicators. **EMPOWER:** build health literacy and sustainable

self-management skills.

FIGURE 8. Holistic Management Framework for Type 2 Diabetes



An eight-pillar, patient-centred framework integrating nutrition, activity, weight management, sleep, stress care, supportive natural approaches, medical care and education.

24. DISCUSSION

Type 2 Diabetes is best understood as a heterogeneous and multifactorial metabolic disease. Different individuals may reach the same diagnosis through different combinations of insulin resistance, beta-cell vulnerability, adiposity, genetics and lifestyle influences. This supports individualized prevention and management rather than simplistic single-cause explanations.

THE BETTER QUESTION IS NOT “WHAT SINGLE FOOD CAUSED DIABETES?” BUT “WHAT COMBINATION OF METABOLIC, GENETIC AND LIFESTYLE FACTORS CONTRIBUTED TO DISEASE DEVELOPMENT?”

25. LIMITATIONS

This paper is an integrative clinical research review and conceptual wellness framework. It does not replace individualized medical diagnosis or treatment, does not prescribe medication, and does not claim that holistic wellness alone cures Type 2 Diabetes. Prevention and management should be individualized with appropriately qualified healthcare professionals.

26. CONCLUSION

Type 2 Diabetes is a progressive disorder of metabolic regulation associated with insulin resistance and progressive beta-cell dysfunction. Important contributors include excess visceral adiposity, physical inactivity, unfavourable

dietary patterns, genetic susceptibility, age, prediabetes, metabolic syndrome, fatty liver disease and selected reproductive or lifestyle-related risk factors.

Understanding these interacting causes creates opportunities for early detection, prevention, risk reduction and better long-term health. The proposed HWRI–T2D Integrated Care Model™ provides a structured pathway integrating medical assessment, metabolic risk identification, lifestyle medicine, nutrition, physical activity, weight management, sleep, psychological wellbeing and patient education.

EARLY DETECTION + RISK REDUCTION + EVIDENCE-BASED CARE + PATIENT EMPOWERMENT = BETTER LONG-TERM METABOLIC HEALTH

27. CLINICAL DISCLAIMER

This publication is intended for educational, academic and public-awareness purposes. Type 2 Diabetes is a serious medical condition requiring appropriate diagnosis, monitoring and treatment. Lifestyle and supportive holistic wellness strategies should not replace professional medical evaluation, prescribed medication or evidence-based diabetes management.

28. KEY SCIENTIFIC REFERENCES

1. World Health Organization. Diabetes Fact Sheet. <https://www.who.int/news-room/fact-sheets/detail/diabetes>
2. National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Symptoms & Causes of Diabetes. <https://www.niddk.nih.gov/health-information/diabetes/overview/symptoms-causes>
3. NIDDK. Insulin Resistance & Prediabetes. <https://www.niddk.nih.gov/health-information/diabetes/overview/what-is-diabetes/prediabetes-insulin-resistance>
4. Centers for Disease Control and Prevention (CDC). Diabetes Risk Factors. <https://www.cdc.gov/diabetes/risk-factors/>
5. CDC. Preventing Type 2 Diabetes. <https://www.cdc.gov/diabetes/prevention-type-2/>
6. American Diabetes Association. Standards of Care in Diabetes—2026. Diagnosis and Classification of Diabetes.
7. American Diabetes Association. Standards of Care in Diabetes—2026. Prevention or Delay of Diabetes and Associated Comorbidities.

SR VAIDYA HOLISTIC WELLNESS RESEARCH INSTITUTE

SRVAIDYA.com | Integrating Ancient Wisdom for Modern Living